

DL2000 Ladder

产品组成

产品名称	产品规格	Cat. No.
DL2000 Ladder	250 μ l	MD1006
DL2000 Ladder	250 μ l $\times$ 5	MD1106

产品储存与有效期

产品可在常温（0-30℃）储存至三年以上。如果长期不用，为防止水分蒸发请于 - 20℃ 储存。


技术支持

杭州新景生物试剂开发有限公司研发部：e-mail: technical@simgen.cn, 电话：400-0099-857。

产品介绍

DL2000 Ladder 由 6 种长度在 100 bp 至 2,000 bp 的 DNA 片段组成，溶解于 1 $\times$ Loading Buffer 中，使用时可取 5-10 μ l 直接电泳，使用非常方便。

特别添加的红色和黄色两种电泳指示染料，不会削弱 DNA 在紫外线下的显色效果，较常用的电泳指示染料（溴酚蓝、二甲苯青等）具有更佳的使用效果。



注意事项

1. 电泳时的加样孔宽度小于 5 mm 时，每次取 5 μ l DNA Ladder 电泳便可得到清晰条带。如果加样孔增宽，须适当增加 DNA Ladder 的加样量。
2. 对 DNA 电泳而言，Agarose 的纯度对 DNA 条带的清晰度影响很大。因此，电泳时应尽量选用质量好的 Agarose，推荐使用胶浓度为 2%~2.5%。
3. 进行 Agarose 电泳时，Agarose 浓度越大，对短片段 DNA 分离性能越好；反之，Agarose 浓度越小，越有利于长片段 DNA 的分离。

DL2000 Ladder

PRODUCT FORMATION

Components	Specification	Cat. No.
DL2000 Ladder	250 μ l	MD1006
DL2000 Ladder	250 μ l $\times$ 5	MD1106

STORAGE

The product can be stored at room temperature (0-30 $^{\circ}$ C) for more than three years. If the product is not used for a long period of time, please store at -20 $^{\circ}$ C to prevent the evaporation of water.

TECHNICAL SUPPORT


TEL: 400-0099-857

E-MAIL: technical@simgen.cn

INTRODUCTION

DL2000 Ladder is composed of 6 individual DNA fragments, presenting 2k, 1k, 750, 500, 250, 100 bp sharp bands respectively. DL2000 DNA Ladder contains 1 $\times$ Loading Buffer, users can apply 5 - 10 μ l in agarose gel electrophoresis directly.

The red and yellow tracking dye in DL2000 Ladder will not weakened the DNA bands under UV light, better than bromophenol blue and xylene cyanol FF.



PRECAUTION

1. Clear bands can be obtained by applying 5 μ l DNA Ladder when the lane width is less than 5 mm. If the lane is wider, loading volume of DNA Ladder should be increased appropriately.
2. For DNA electrophoresis, agarose purity is of great significance to DNA band definition. Therefore, agarose with good quality should be used and gel concentration of 2%~2.5% is recommended.
3. During agarose electrophoresis, the concentration of agarose is closely associated with the separation of DNA fragments. High agarose concentration is ideal for the separation of the short DNA fragments. While low agarose concentration is ideal to separate the long DNA fragments.